

## First record of the Mourning Gecko, *Lepidodactylus lugubris* (Duméril & Bibron), from Bonaire

Gerard van Buurt<sup>1,\*</sup> and JCGJ Smulders<sup>2</sup>

<sup>1</sup>Kaya Oy Sprock 18, Curaçao

<sup>2</sup>Schoolstraat 32, 6093EH Heythuysen, The Netherlands

\*Corresponding author (gvanbuurt@gmail.com)

**Edited by:** S. Blair Hedges. **Date of publication:** 3 February 2022.

**Citation:** van Buurt G, Smulders JCGI (2022) First record of the Mourning Gecko, *Lepidodactylus lugubris* (Duméril & Bibron), from Bonaire. *Caribbean Herpetology*, 82, 1.

**DOI:** <https://doi.org/10.31611/ch.82>

On the 14th of December, 2020, an individual of *Lepidodactylus lugubris* was found and photographed in Bonaire (Dutch Caribbean) by Hans Smulders in a garden at the following address: Sabadeco Shores 28 (12.198174, -68.302846), 16 m elevation. The animal was found around some potted plants. This species has not been reported previously from Bonaire (Behm *et al.* 2019).



**Figure 1.** *Lepidodactylus lugubris* from Bonaire. Photo by JCGJ Smulders.

### References

Behm JE, van Buurt G, DiMarco BM, Eilers J, Irian CG, Langhans KE, McGrath K, Tran TJ, Helmus MR (2019) First records of the mourning gecko (*Lepidodactylus lugubris* Duméril and Bibron, 1836), common house gecko (*Hemidactylus frenatus* in Duméril, 1836), and Tokay gecko (*Gekko gekko* Linnaeus, 1758) on Curaçao, Dutch Antilles, and remarks on their Caribbean distributions. *BioInvasions Records* 8, 34–44. **Article**